

Evidence Synthesis / Meta-analysis

Session 3, Lecture 6: Small study effects

Guido Knapp¹, Gerta Rücker², Guido Schwarzer²

¹ Department of Statistics, TU Dortmund University

² Center for Medical Biometry and Medical Informatics, University of Freiburg

ruecker@imbi.uni-freiburg.de

DAGStat 2016, Göttingen, 14 March 2016



Overview Lecture 6

- ▶ Types of bias
 - ▶ Publication bias and related biases
 - ▶ Small-study effects
- ▶ Diagnosis of small-study effects
 - ▶ Funnel plot
 - ▶ Funnel plot tests
- ▶ Adjustment for small-study effects
 - ▶ Trim and fill method
 - ▶ Copas selection model (short description of method)
 - ▶ Adjustment by regression

Bias in meta-analyses: Small-study effects

- ▶ **Publication bias** (Easterbrook et al., 1991; Rothstein et al., 2005): Small studies tend to be published only if they show a large effect
- ▶ Related types of bias: Studies having 'significant' results tend to be
 - ▶ published in high-ranking English language journals (*Language bias*) (Egger et al., 1997b)
 - ▶ published faster than studies without a 'significant' result (*Time lag bias*) (Higgins and Green, 2009)
 - ▶ published more than once (*Multiple publication bias*) (Gøtzsche, 1989)
 - ▶ cited more often than studies without a 'significant' result, and therefore are more easily detectable in literature searches (*Citation bias*) (Nieminen et al., 2007)

Small-study effects

Smaller trials show different, often larger, treatment effects than large ones (Sterling et al., 1995; Sterne et al., 2000; Rothstein et al., 2005)


- ▶ Potential causes of small-study effects
 - ▶ **Publication bias**: Small studies tend to be published preferably if they show a large effect (Easterbrook et al., 1991)
 - ▶ **Selective outcome reporting bias**: Studies present selected outcomes (Chan et al., 2004a,b; Williamson and Gamble, 2005)
 - ▶ **Selective analysis reporting bias**: Studies choose a method of analysis that leads to larger effects (Ioannidis et al., 2014)
 - ▶ **Clinical heterogeneity** between patients in large and small trials
 - ▶ **For binary data: Statistical correlation** between treatment effect estimate and its variance (Schwarzer et al., 2002)
 - ▶ **Coincidence**
- ▶ Graphical representation of small-study effects
 - ▶ Funnel plot (Light and Pillemer, 1984; Sterne and Egger, 2001)


Funnel plot


Horizontal axis: (log) treatment effect


Vertical axis: a measure of precision; various versions in the literature:

- ▶ Sample size
- ▶ Inverse variance
- ▶ Standard error on a reversed axis (preferred, Sterne and Egger (2001))
 - ▶ confidence intervals increasing linearly
 - ▶ sufficient space for imprecise (small) studies (particularly interesting for diagnosis of small-study effects)











Bias in meta-analysis \Leftrightarrow Asymmetry in funnel plot



Bias in meta-analysis \Leftrightarrow Asymmetry in funnel plot



Funnel plot

Example: NSAIDS data


Meta-analysis of 37 placebo-controlled randomized trials on the effectiveness and safety of topical non-steroidal anti-inflammatory drugs (NSAIDS) in acute pain (Moore et al., 1998)

Part of R package **metasens**


```
# Perform meta-analysis  
ms1 <- metabin(Ee, Ne, Ec, Nc, data = nsaid, sm = "OR")
```

```
# Create funnel plot  
funnel(ms1)
```

Funnel plot of NSAIDS data



Contour-enhanced funnel plot of NSAIDs data



Funnel plot-based tests for small-study effects

- ▶ **Idea:** Test for asymmetry in the funnel plot as an indication for bias
- ▶ **Method:** Test for association between treatment effect and standard error
- ▶ **Assumption:** No association between treatment effect and standard error (or precision) if there is no small-study effect
- ▶ **Limitation:** Strictly valid only for normally distributed outcomes
- ▶ Criticised by some authors (Terrin et al., 2005; Lau et al., 2006)

Funnel plot tests for asymmetry: Overview

Rank correlation tests (not considered here)

- ▶ Test by Begg and Mazumdar (1994)
 - ▶ Modification for binary data: Test by Schwarzer et al. (2007)

Regression tests

- ▶ Test by Egger et al. (1997a)
- ▶ Modifications for binary data
 - ▶ Test by Harbord et al. (2006)
 - ▶ Test by Macaskill et al. (2001)
 - ▶ Test by Peters et al. (2006)
 - ▶ Arcsine test (Rücker et al., 2008)

Regression tests: Basic idea

Choose an effect measure, say, the mean difference

Null-hypothesis ('*No small study effects*'): Treatment effect does not depend on precision

1. Regress the treatment effect on the standard error, using inverse variance weights
2. Test null-hypothesis of zero slope

Often called Egger's test (Egger et al., 1997a)

Note: Strictly valid only for continuous data (data normally distributed)!

Nevertheless often applied to binary data, preferably in a modified version

Egger's test and its modifications for binary data in R

- ▶ Egger's test (Egger et al., 1997a)
 - ▶ Use R: `metabias(ms1, method = "linreg")`
- ▶ Harbord's score test (Harbord et al., 2006)
 - ▶ Uses a score-based estimate for the odds ratio
 - ▶ Advantage: Variance estimate depends only on marginal totals
 - ▶ Use R: `metabias(ms1, method = "score")`
- ▶ Peters' test (Peters et al., 2006)
 - ▶ Uses the usual odds ratio estimate and $1/n$ as regressor
 - ▶ Advantage: Study weights depending only on marginal totals
 - ▶ Use R: `metabias(ms1, method = "peters")`
- ▶ Arcsine test (Rücker et al., 2008)
 - ▶ Uses the arcsine difference instead of the odds ratio
 - ▶ Advantage: Variance depends only on group sample sizes
 - ▶ Use R: `ms1.asd <- update(ms1, sm = "ASD")`
`metabias(ms1.asd, method = "linreg")`

Recommendations on testing for funnel plot asymmetry

Sterne et al. (2011), BMJ:

- ▶ Funnel plot tests only when there are at least 10 studies (rule of thumb; argument `k.min` in function `metabias`)
- ▶ Recommendation for continuous outcomes:
Linear regression test (Egger et al., 1997a)
- ▶ For binary outcomes:
Use one of the modifications of Egger's test (Harbord et al., 2006; Peters et al., 2006; Rücker et al., 2008)
- ▶ Bias cannot be excluded if test for funnel plot asymmetry is non-significant
- ▶ Test performance deteriorates if between-study heterogeneity increases

Adjusting for small-study effects

Three approaches

- ▶ **Trim and fill method**
(Duval and Tweedie, 2000a,b)
- ▶ **Copas selection model for publication bias**
(Copas, 1999; Copas and Shi, 2000, 2001)
- ▶ **Adjustment by regression**
(Copas and Malley, 2008; Stanley, 2008; Moreno et al., 2009a,b; Rücker et al., 2010)


Trim and fill method

1. Estimate the number of studies in the outlying part of the funnel plot using rank-based methods;
2. remove (**trim**) these studies and do meta-analysis on the remaining studies;
3. consider the estimate from the 'trimmed' meta-analysis as the true center of the funnel;
4. for each 'trimmed' study, create (**fill**) an additional study as the mirror image about the center of funnel plot;
5. do meta-analysis on original and filled studies.

```
# Perform Trim and fill analysis  
tf1 <- trimfill(ms1)
```

```
# Create funnel plot including filled-in studies  
funnel(tf1)
```

Trim and fill plot of NSAIDS data



Trim and fill method

```
# Print results of Trim and fill analysis
```

```
print(tfl, digits = 2)
```

```
##              OR          95%-CI %W(random)
## 1          6.57 [2.11; 20.48]      2.13
*** Output truncated ***
## 37          5.69 [1.51; 21.42]      1.91
## Filled: 37    0.95 [0.25; 3.56]      1.91
## Filled: 27    0.86 [0.16; 4.68]      1.53
## Filled: 16    0.82 [0.25; 2.73]      2.05
*** Output truncated ***
## Filled: 32    0.05 [0.00; 1.08]      0.68
##
## Number of studies combined: k=51 (with 14 added studies)
##
##              OR          95%-CI z  p-value
## Random effects model 2.45 [1.83; 3.28] 6 < 0.0001
##
## Quantifying heterogeneity:
## tau^2 = 0.7113; H = 1.93 [1.68; 2.22]; I^2 = 73.2% [64.7%; 79.7%]
*** Output truncated ***
```

Copas selection model

Combine two models:

1. Usual random effects model for **treatment effect**
2. A model for the **selection process** with a parameter controlling how chance of publication depends on precision $1/s_k$ (where s_k is the within-study standard error)

Selection/publication bias is modelled by a parameter representing the **correlation** between effect size and selection probability

- ▶ Implemented in function **copas** of R package **metasens** (earlier: **copas**) (Carpenter et al., 2009a)
- ▶ Sensitivity analysis necessary
- ▶ Details given in (Copas, 1999; Copas and Shi, 2001)

Adjustment by regression (Rücker et al., 2010)

► Random effects model

$$\hat{\theta}_k = \theta_k + \sigma_k \eta_k, \quad \eta_k \sim N(0, 1)$$

$$\theta_k = \theta + \tau \delta_k, \quad \delta_k \sim N(0, 1)$$

- $\hat{\theta}_k$ observed treatment effect in study k ($k = 1, \dots, K$)
- θ_k true treatment effect in study k
- θ overall treatment effect
- σ_k^2 within-study sampling variance, τ^2 between-study variance

► Equivalent:

$$\hat{\theta}_k = \theta + \sqrt{\sigma_k^2 + \tau^2} \epsilon_k, \quad \epsilon_k \sim N(0, 1)$$

Adjustment by regression

- ▶ Random effects model:

$$\hat{\theta}_k = \theta + \sqrt{\sigma_k^2 + \tau^2} \epsilon_k, \quad \epsilon_k \sim N(0, 1)$$

- ▶ **Extended random effects model** taking account of possible small study effects by allowing the effect to depend on the standard error:

$$\hat{\theta}_k = \theta + \sqrt{\sigma_k^2 + \tau^2} (\alpha + \epsilon_k), \quad \epsilon_k \sim N(0, 1)$$

Additional parameter α represents bias introduced by small-study effects ('publication bias')

Adjustment by regression

- ▶ **Extended random effects model**, taking account of possible small study effects by allowing the effect to depend on the standard error:

$$\hat{\theta}_k = \theta + \sqrt{\sigma_k^2 + \tau^2} (\alpha + \epsilon_k), \quad \epsilon_k \sim N(0, 1)$$

- ▶ **Additional parameter α** represents bias introduced by small-study effects ('publication bias')
 - ▶ For a very small study k , we have $\sigma_k^2 \rightarrow \infty$ and therefore

$$E \left(\frac{\hat{\theta}_k - \theta}{\sigma_k} \right) \rightarrow \alpha \text{ Small study bias}$$

- ▶ For a very large study k , we have $\sigma_k^2 \rightarrow 0$ and therefore

$$E(\hat{\theta}_k) \rightarrow \theta + \tau \alpha \text{ Adjusted effect of large study}$$


- ▶ Implemented in function **limitmeta** of R package **metasens** (Carpenter et al., 2009a)

Adjustment by regression

```
# Perform limit meta-analysis  
l1 <- limitmeta(ms1)
```

```
# Create funnel plot with adjusted regression line  
funnel(l1, col.line = "red", lwd.line = 2)
```

Funnel plot with adjusted regression line for NSAIDS data



Adjustment by regression

```
# Print results of regression adjustment (limit meta-analysis)
print(summary(l1), digits = 2)

## Result of limit meta-analysis:
##
## Random effects model   OR      95%-CI    z      pval
## Adjusted estimate 1.84 [1.26; 2.68] 3.17  0.0015
## Unadjusted estimate 3.73 [2.80; 4.97] 9.01 < 0.0001
##
## Quantifying heterogeneity:
## tau^2 = 0.4670; I^2 = 68.3% [55.5%; 77.4%]; G^2 = 91.5%
##
## Test of heterogeneity:
##      Q d.f.  p.value
## 113.52   36 < 0.0001
##
## Test of small-study effects:
##      Q-Q' d.f.  p.value
##  44.20     1 < 0.0001
##
## Test of residual heterogeneity beyond small-study effects:
##      Q' d.f.  p.value
##  69.32    35  0.0005
```

Compare estimates for NSAIDS example

Model	Odds ratio [95% CI]
Fixed effect model	2.89 [2.49; 3.35]
Random effects model	3.73 [2.80; 4.97]
Trim and fill (random effects estimate)	2.45 [1.83; 3.28]
Copas selection model	1.82 [1.46; 2.26]
Regression adjustment	1.84 [1.26; 2.68]

Adjusting for small-study effects: Summary

Three approaches

- ▶ Trim and fill method
 - ▶ R function **trimfill** in R package **meta**
 - ▶ Not model-based, somewhat ad hoc
- ▶ Copas selection model for publication bias
 - ▶ R function **copas**, implemented in R package **metasens**
 - ▶ Model-based, needs sensitivity analysis
 - ▶ Sometimes associated with estimation problems (Carpenter et al., 2009b)
- ▶ Adjustment by regression
 - ▶ R function **limitmeta**, implemented in R package **metasens**
 - ▶ Model-based, extension of the regression test

References

- Begg, C. B. and Mazumdar, M. (1994). Operating characteristics of a rank correlation test for publication bias. *Biometrics*, 50:1088–1101.
- Carpenter, J., Rücker, G., and Schwarzer, G. (2009a). copas: An R package for fitting the Copas selection model. *The R Journal*, 1(2):31–36.
- Carpenter, J. R., Schwarzer, G., Rücker, G., and Küntzler, R. (2009b). Empirical evaluation showed that the Copas selection model provided a useful summary in 80% of meta-analyses. *Journal of Clinical Epidemiology*, 62:624–631.
- Chan, A.-W., Hróbartsson, A., Haahr, M. T., Gøtzsche, P. C., and Altman, D. G. (2004a). Empirical evidence for selective reporting of outcomes in randomized trials. comparison of protocols to published articles. *Journal of the American Medical Association*, 291:2457–2465.
- Chan, A.-W., Kroleza-Jeric, K., Schmid, I., and Altman, D. G. (2004b). Outcome reporting bias in randomized trials funded by the canadian institutes of health research. *Canadian Medical Association Journal*, 171:735–740.
- Copas, J. (1999). What works?: Selectivity models and meta-analysis. *Journal of the Royal Statistical Society, Series A*, 162:95–109.
- Copas, J. and Shi, J. Q. (2000). Meta-analysis, funnel plots and sensitivity analysis. *Biostatistics*, 1:247–262.
- Copas, J. B. and Malley, P. F. (2008). A robust p-value for treatment effect in meta-analysis with publication bias. *Statistics in Medicine*, 27(21):4267–4278.

- Copas, J. B. and Shi, J. Q. (2001). A sensitivity analysis for publication bias in systematic reviews. *Statistical Methods in Medical Research*, 10:251–265.
- Duval, S. and Tweedie, R. (2000a). A nonparametric “Trim and Fill” method of accounting for publication bias in meta-analysis. *Journal of the American Statistical Association*, 95:89–98.
- Duval, S. and Tweedie, R. (2000b). Trim and Fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56:455–463.
- Easterbrook, P. J., Berlin, J. A., Gopalan, R., and Matthews, D. R. (1991). Publication bias in clinical research. *The Lancet*, 337:867–872.
- Egger, M., Smith, G. D., Schneider, M., and Minder, C. (1997a). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315:629–634.
- Egger, M., Zellweger-Zähner, T., Schneider, M., Junker, C., Lengeler, C., and Antes, G. (1997b). Language bias in randomised controlled trials published in English and German. *The Lancet*, 350:326–329.
- Gøtzsche, P. C. (1989). Multiple publication of reports of drug trials. *European Journal of Clinical Pharmacology*, 36:429–432.
- Harbord, R. M., Egger, M., and Sterne, J. A. (2006). A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in Medicine*, 25(20):3443–3457.
- Higgins, J. P. and Green, S. (2009). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.0.2*. Chichester.

- Ioannidis, J., Munafo, M., Fusar-Poli, P., Nosek, B., and David, S. (2014). Publication and other reporting biases in cognitive sciences: detection, prevalence, and prevention. *Trends in Cognitive Sciences*, 18(5):235–241. doi: 10.1016/j.tics.2014.02.010.
- Lau, J., Ioannidis, J. P. A., Terrin, N., Schmid, C. H., and Olkin, I. (2006). The case of the misleading funnel plot. *British Medical Journal*, 333:597–600.
- Light, R. J. and Pillemer, D. B. (1984). *Summing up. The science of reviewing research*. Harvard University Press, Cambridge, Massachusetts.
- Macaskill, P., Walter, S. D., and Irwig, L. (2001). A comparison of methods to detect publication bias in meta-analysis. *Statistics in Medicine*, 20:641–654.
- Moore, R. A., Tramer, M. R., Carroll, D., Wiffen, P. J., and McQuay, H. J. (1998). Quantitative systematic review of topically applied non-steroidal anti-inflammatory drugs. *British Medical Journal*, 316(7128):333–338.
- Moreno, S., Sutton, A., Ades, A., Stanley, T., Abrams, K., Peters, J., and Cooper, N. (2009a). Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study. *BMC Medical Research Methodology*, 9:2.
- Moreno, S. G., Sutton, A. J., Turner, E. H., Abrams, K. R., Cooper, N. J., Palmer, T. P., and Ades, A. E. (2009b). Novel methods to deal with publication biases: secondary analysis of antidepressant trials in the FDA trial registry database and related journal publications. *BMJ*, 339. doi: 10.1136/bmj.b2981.
- Nieminen, P., Rücker, G., Miettinen, J., Carpenter, J. R., and Schumacher, M. (2007). Statistically significant papers in psychiatry were cited more often than others. *Journal of Clinical Epidemiology*, 60(9):939–946.

- Peters, J. L., Sutton, A. J., Jones, D. R., Abrams, K. R., and Rushton, L. (2006). Comparison of two methods to detect publication bias in meta-analysis. *Journal of the American Medical Association*, 295:676–680.
- Rothstein, H. R., Sutton, A. J., and Borenstein, M. (2005). *Publication bias in meta analysis: prevention, assessment and adjustments*. Wiley, Chichester.
- Rücker, G., Schwarzer, G., Carpenter, J., Binder, H., and Schumacher, M. (2010). Treatment effect estimates adjusted for small-study effects via a limit meta-analysis. *Biostatistics*, 12(1):122–142. Doi:10.1136/jme.2008.024521.
- Rücker, G., Schwarzer, G., and Carpenter, J. R. (2008). Arcsine test for publication bias in meta-analyses with binary outcomes. *Statistics in Medicine*, 27(5):746–763.
- Schwarzer, G., Antes, G., and Schumacher, M. (2002). Inflation of type I error rate in two statistical tests for the detection of publication bias in meta-analyses with binary outcomes. *Statistics in Medicine*, 21:2465–2477.
- Schwarzer, G., Antes, G., and Schumacher, M. (2007). A test for publication bias in meta-analysis with sparse binary data. *Statistics in Medicine*, 26:721–733.
- Schwarzer, G., Carpenter, J. R., and Rücker, G. (2010). Empirical evaluation suggests Copas selection model preferable to trim-and-fill method for selection bias in meta-analysis. *Journal of Clinical Epidemiology*, 63:282–288.
- Stanley, T. D. (2008). Meta-regression methods for detecting and estimating empirical effects in the presence of publication selection. *Oxford Bulletin of Economics and Statistics*, 70:105–127.

- Sterling, T. D., Rosenbaum, W. L., and Weinkam, J. J. (1995). Publication decisions revisited: The effect of the outcome of statistical tests on the decision to publish and vice versa. *The American Statistician*, 49:108–112.
- Sterne, J. A. C. and Egger, M. (2001). Funnel plots for detecting bias in meta-analysis: Guideline on choice of axis. *Journal of Clinical Epidemiology*, 54:1046–1055.
- Sterne, J. A. C., Gavaghan, D., and Egger, M. (2000). Publication and related bias in meta-analysis: Power of statistical tests and prevalence in the literature. *Journal of Clinical Epidemiology*, 53:1119–1129.
- Sterne, J. A. C., Sutton, A. J., Ioannidis, J. P. A., Terrin, N., Jones, D. R., Lau, J., Carpenter, J., Rücker, G., Harbord, R. M., Schmid, C. H., Tetzlaff, J., Deeks, J. J., Peters, J., Macaskill, P., Schwarzer, G., Duval, S., Altman, D. G., Moher, D., and Higgins, J. P. T. (2011). Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *British Medical Journal*, 343:d4002. doi: 10.1136/bmj.d4002.
- Terrin, N., Schmid, C. H., and Lau, J. (2005). In an empirical evaluation of the funnel plot, researchers could not visually identify publication bias. *Journal of Clinical Epidemiology*, 58(9):894–901.
- Williamson, P. R. and Gamble, C. (2005). Identification and impact of outcome selection bias in meta-analysis. *Statistics in Medicine*, 24(10):1547–1561.

Binary outcome: Notation

Result of study k

Event	yes	no	Total
Treatment	a_k	b_k	$a_k + b_k$
Control	c_k	d_k	$c_k + d_k$
Total	$a_k + c_k$	$b_k + d_k$	n_k

Modification I: Harbord's score test

- ▶ Take the efficient score

$$Z_k = a_k - (a_k + b_k)(a_k + c_k)/n_k$$

of the log odds ratio with variance (Fisher's information)

$$V_k = (a_k + b_k)(a_k + c_k)(b_k + d_k)(c_k + d_k)/[n_k^2(n_k - 1)]$$

- ▶ Z_k/V_k is an estimate for the log odds ratio, if the true value of θ is not too far from 0 (standard likelihood theory)
- ▶
 1. Regress treatment effect estimate Z/V (with variance $1/V$) on its standard error $1/\sqrt{V}$ with study weights V
 2. Test null-hypothesis of zero slope
- ▶ Advantage: Variance estimate depends only on marginal totals

Modification II: Macaskill's/Peters' test

- ▶ Choose as effect measure the log odds ratio
- ▶ Let $\hat{\theta}_k$ be the effect estimate of study k , $a_k + c_k$ the number of events, $b_k + d_k$ the number of non-events in study k
 1. Regress treatment effect $\hat{\theta}$ on sample size n (Macaskill) or inverse sample size $1/n$ (Peters) with study weights $(1/(a + c) + 1/(b + d))^{-1}$ (both)
 2. Test null-hypothesis of zero slope
- ▶ Advantage: Variance estimate depends only on marginal totals

Modification III: Arcsine test

- ▶ With

$$p_{T,i} = \frac{a_k}{a_k + b_k} \quad \text{and} \quad p_{C,i} = \frac{c_k}{c_k + d_k}$$

choose as effect measure the arcsine difference

$$\Delta_k = \arcsin \sqrt{p_{T,i}} - \arcsin \sqrt{p_{C,i}}$$

- ▶ 1. Regress treatment effect Δ on standard error $\sqrt{\frac{1}{a+b} + \frac{1}{c+d}}$
with weights $1/(\frac{1}{a+b} + \frac{1}{c+d})$
- 2. Test null-hypothesis of zero slope
- ▶ Advantage: Variance estimate independent of mean treatment effect

Overview: Regression tests

Test	Measure	Regressor	Weights
Egger's test	$\hat{\theta}$	$\text{SE}(\hat{\theta})$	$1/\widehat{\text{Var}}(\hat{\theta})$
Harbord's score test	Z/V	$1/\sqrt{V}$	V
Peters' test	$\hat{\theta}$	$1/n$	$1/(\frac{1}{a+c} + \frac{1}{b+d})$
Arcsine test	Δ	$\sqrt{\frac{1}{a+b} + \frac{1}{c+d}}$	$1/(\frac{1}{a+b} + \frac{1}{c+d})$

Compare results of fixed effect and random effects model

```
summary(ms1)
```

```
## Number of studies combined: k=37
##
##                               OR           95%-CI           z   p-value
## Fixed effect model    2.9809 [2.5854; 3.4368] 15.0409 < 0.0001
## Random effects model 3.7345 [2.8039; 4.9740]  9.0105 < 0.0001
##
## Quantifying heterogeneity:
## tau^2 = 0.4670; H = 1.78 [1.5; 2.1]; I^2 = 68.3% [55.5%; 77.4%]
##
## Test of heterogeneity:
##      Q d.f.  p-value
## 113.52   36 < 0.0001
##
## Details on meta-analytical method:
## - Mantel-Haenszel method
## - DerSimonian-Laird estimator for tau^2
## - Continuity correction of 0.5 in studies with zero cell frequencies
```

Funnel plot and contour-enhanced funnel plot

```
# Load the data  
data(nsaid)
```

```
# Perform meta-analysis  
ms1 <- metabin(Ee, Ne, Ec, Nc, data = nsaid, sm = "OR")
```

```
# Create funnel plot  
funnel(ms1)
```

```
# Create contour-enhanced funnel plot  
funnel(ms1, comb.random = FALSE, pch = 16,  
        contour = c(0.9, 0.95, 0.99),  
        col.contour = c("green", "yellow", "pink"))  
legend(0.25, 1.25,  
       c("0.1 > p > 0.05", "0.05 > p > 0.01", "< 0.01"),  
       fill = c("green", "yellow", "pink"), bty = "n")
```

Copas selection model

Combine two models:

1. Usual random effects model for **treatment effect**
2. A model for the **selection process** with a parameter controlling how chance of publication depends on precision $1/s_k$ (where s_k is the within-study standard error)





Selection/publication bias is modelled by a parameter representing the **correlation** between effect size and selection probability

- ▶ Implemented in function **copas** of R package **metasens** (earlier: **copas**) (Carpenter et al., 2009a)
- ▶ Sensitivity analysis necessary

```
# Use Copas selection model  
c1 <- copas(ms1)
```

```
# Create plots for sensitivity analysis  
plot(c1)
```

Copas selection model plot of NSAIDS data



Copas selection model

```
# Print results of Copas selection model
print(summary(c1), digits = 2)


## Summary of Copas selection model analysis:
##
##          publprob   OR      95%-CI pval.treat pval.rsb N.unpubl
##          1.00 3.73 [2.77; 5.02] < 0.0001 < 0.0001      0
##          0.98 3.67 [2.89; 4.65] < 0.0001 < 0.0001      0
##          0.84 3.32 [2.43; 4.55] < 0.0001 < 0.0001      3
##          0.70 3.01 [2.21; 4.09] < 0.0001 < 0.0001      8
##          0.59 2.72 [2.01; 3.69] < 0.0001 0.0004     14
##          0.50 2.46 [1.82; 3.34] < 0.0001 0.0014     20
##          0.43 2.23 [1.65; 3.02] < 0.0001 0.0054     28
##
##      Copas model (adj)
## Random effects model 3.73 [2.80; 4.97] < 0.0001
##
## Significance level for test of residual selection bias: 0.1
##
## Legend:
## publprob   - Probability of publishing study with largest standard error
## pval.treat  - P-value for hypothesis of overall treatment effect
## pval.rsb    - P-value for hypothesis that no selection remains unexplained
```

Copas selection model: Change sensitivity parameters

```
# Use Copas selection model, range of (gamma0, gamma1) modified  
c2 <- copas(ms1, gamma0.range = c(-1, 2), gamma1.range = c(0, 1))
```

```
# Create plots for sensitivity analysis  
plot(c2)
```


Copas selection model plot of NSAIDS data



Copas selection model

```
# Print results of Copas selection model
```

```
print(summary(c2), digits = 2)
```

```
## Summary of Copas selection model analysis:
```

```
##
```

	publprob	OR	95%-CI	pval.treat	pval.rsb	N.unpubl
##	1.00	3.73	[2.77; 5.02]	< 0.0001	< 0.0001	0
##	0.75	3.33	[2.45; 4.52]	< 0.0001	< 0.0001	4
##	0.49	2.73	[2.05; 3.64]	< 0.0001	0.0018	15
##	0.35	2.24	[1.72; 2.92]	< 0.0001	0.0356	30
##	0.26	1.82	[1.46; 2.26]	< 0.0001	0.3273	48

```
##
```

##	Copas model (adj)	1.82	[1.46; 2.26]	< 0.0001	0.3273	48
##	Random effects model	3.73	[2.80; 4.97]	< 0.0001		

```
##
```

```
## Significance level for test of residual selection bias: 0.1
```

```
##
```

```
## Legend:
```

```
## publprob - Probability of publishing study with largest standard error
## pval.treat - P-value for hypothesis of overall treatment effect
## pval.rsb - P-value for hypothesis that no selection remains unexplained
## N.unpubl - Approximate number of unpublished studies suggested by model
```